



Acacia auriculiformis

Jøker, Dorthé

Published in:
Seed Leaflet

Publication date:
2000

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Jøker, D. (2000). *Acacia auriculiformis*. *Seed Leaflet*, (2).

SEED LEAFLET

No. 2 March 2000



Acacia auriculiformis Cunn. ex Benth.

Taxonomy and nomenclature

Family: Fabaceae (Mimosoideae)

Synonyms: *Acacia auriculaeformis* A. Cunn. ex Benth., *Racosperma auriculiforme* (A. Cunn. ex Benth.) Pedley.

Vernacular/common names: Northern black wattle (Australian trade name); coast wattle, ear pod wattle.

Distribution and habitat

Native to Australia, Papua New Guinea and Indonesia in hot humid and sub-humid lowlands with mean annual rainfall of 800-2500 mm and mean annual temperature of 20-30°C. Often found on river banks and in coastal areas.

It is cultivated widely in the tropics within an altitude range of 0-500 (-1000) masl, and even though frost does not occur in its natural range, it tolerates light frost. It is exceptionally tolerant to soil type in regard to fertility, salinity and pH. It can grow on acid mine spoils with pH 3 and on alkaline beach sands with pH 8-9. It does not tolerate shade or strong winds.


Isoenzyme analysis has revealed a marked genetic variation with 3 distinct groups corresponding to the geographic distribution in Papua New Guinea, Queensland and Northern Territory. Field trials show that provenances from Papua New Guinea have the highest production while those from Queensland have a high proportion of single stems. Those from the Northern Territory are inferior in both growth and form. The hybrid *A. auriculiformis* x *A. mangium* has shown desirable characteristics.

Uses

This species can grow under even the most difficult conditions in the tropics. The rapid early growth rate, ability to fix nitrogen, tolerance of infertile, acid, alkaline, saline or seasonally waterlogged soils and moderately dry seasons makes it very suitable for rehabilitation of degraded land. A valuable feature is the ability to compete with *Imperata cylindrica* reducing the grass to a sparse ground cover. The wood is excellent for firewood, charcoal, paper pulp and lighter construction work. Because of its spreading and competitive surface rooting habit it is not widely used in agroforestry systems but intercropping with peanut, rice, mung beans and kenaf has been successful.

Botanical description

A large shrub or medium-sized, evergreen tree, usually 8-20 m tall, on good sites up to 35 m. Bark grey or brown, longitudinally fissured. Leaves (phyllodes) 8-20 cm long, glabrous and curved, with 3 prominent nerves (four in *A. mangium*). Flowers bisexual, creamy yellow, scented, in up to 8.5 cm long spikes.



Seed with funicle, flowering branch and pod. Copyright: CSIRO, Forestry & Forest Products, Australian Tree Seed Centre.

Fruit and seed description

Fruit: flat, dehiscent, somewhat woody pod, 6.5 cm long, 1.5 cm wide, strongly curved and with undulate margins.

Seed: shiny black or brown, encircled by a long, red or yellow funicle. There are 55,000-75,000 seeds/kg.

Flowering and fruiting habit

The yellow flower spikes can be found on individual trees throughout the year but there is usually a distinct peak flowering season which may vary considerably with location.

Pollination is carried out by a wide range of insects. Seed is produced at an early age and normally in large quantities.

Phenology data:

Locality	Flowering	Fruiting
N. Territory	Apr-June	Aug-Oct
Queensland	Feb-May	Oct-Apr
Kuala Lumpur	Feb-May	Oct-Apr
Sabah	Feb-May	Oct-Apr
Java	Mar-June	
Thailand	Aug-Feb	
Vietnam	June-July	Nov-Mar

Harvest

Collection from the tree or from the ground. If collection is too early, it can be difficult to open the pods while late collection may result in loss of seeds.

Processing and handling

The pods are dried in the sun until they open. After extraction the funicle can be removed manually by rubbing the seed over a sieve.

Storage and viability

Seeds are orthodox and can be stored for several years in closed plastic bags. Moisture content for storage should be about 7%.

Dormancy and pretreatment

Pretreatment by scarification or immersion in boiling water for 1 min and left to soak for 24 hours.

Sowing and germination

Germination of 40-100% occurs after 6-15 days. Propagation by container plants, direct sowing or vegetatively by cuttings. Nursery techniques are described in Doran and Turnbull (1997).

Selected readings

Booth, T.H., Turnbull, J.W., 1994. *Domestication of lesser-known tropical tree species: The Australian experience.* In: Leakey RRB, Newton AC, eds. *Tropical trees: The Potential for Domestication and Rebuilding of Forest Resources.* ITE.

Boland, D.J., 1989. *Trees for the tropics. Growing Australian multipurpose trees and shrubs in Developing Countries.* ACIAR Monograph, No. 10, ii + 247 pp.; 11 pp. of ref.

Doran, J.C., Gunn, B.V., 1987. *Treatments to promote seed germination in Australian acacias.* In: Turnbull JW, ed. *Australian Acacias in Developing Countries.* Proceedings of an International Workshop, Gympie, Qld., Australia, 4-7 August 1986. ACIAR Proceedings No 16:57-63

Doran, J.C., Turnbull, J.W., 1997. *Australian trees and shrubs: Species for land rehabilitation and farm planting in the tropics.* ACIAR Monograph No. 24. 384 pp.

Faridah Hanum, I., van der Maesen, L.J.G., eds., 1997. *Plant Resources of South-East Asia. No. 11. Auxillary plants.* Leiden, Netherlands: Backhuys. 389 pp.

Khongsak Pinyopusarerk, 1996. *Acacia auriculiformis - a multipurpose tropical wattle.* FACT Sheet 96-05

Padma, V., Reddy, B.M., Satyanarayana, G., 1993. *Breaking dormancy in certain Acacia spp. by pre-sowing seed treatments.* Seed Research, publ. 1995, 21(1):26-30; 4 ref.

Wickneswari, R. and Norwati, M., 1991. *Genetic structure of natural populations of Acacia auriculiformis in Australia and Papua New Guinea.* In: J.W. Turnbull (ed), *Advances in tropical acacia research.* ACIAR Proceedings No. 35.



Forest tree form. Bensbach River, Balamuk, Western Provenance, Papua New Guinea. Photo: Maurice McDonald, CSIRO, Forestry & Forest Products, Australian Tree Seed Centre.

THIS NOTE WAS PREPARED BY
DANIDA FOREST SEED CENTRE

Author: Dorte Jøker

Danida Forest Seed Centre
Krogerupvej 21
DK-3050 Humlebaek
Denmark

Phone: +45-49190500
Fax: +45-49160258
Email: dfsc@sns.dk
Website: www.dfsc.dk